## Chapter 6 LITERATURE CITED

- Baker, B. 1990. Draft Caloosahatchee water quality based effluent limitations documentation (Lee County). Florida Department of Environmental Regulation, Water Quality Tech. Ser. 2(121)
- Bierman, V. 1993. Performance report for the Caloosahatchee Estuary salinity modeling. SFWMD expert assistance contract deliverable from Limno-Tech, inc.
- Bortone, S.A., M. P. Smith, 1998. Indexed bibliography to the biology and life history of tape grass (*Vallisneria americana*). Florida Center for Environmental Studies. Technical Series No. 4.
- Bortone, S.A. and Turpin, R.K. 1999. Tapegrass Life History Metrics Associated with Environmental Variables in a Controlled Estuary. Report to South Florida Water Management District, West Palm Beach, FL.
- Bourn, W.S. 1932. Ecological and physiological studies on certain aquatic angiosperms. Contrib. Boyce Thompson Inst. 4: 425 496.
- Bourn, W.S. 1934. Sea-water tolerance of *Vallisneria spiralis* L. and *Potomogeton foliosus* Raf. Contrib. Boyce Thompson Inst. 6: 303 308.
- Carter, V., J. W. Barko, G. L. Godshalk, and N. B. Rybicki. 1988. Effects of submersed macrophytes on water quality in the tidal Potomac River, Maryland. Journal of Freshwater Ecology 4:493-501.\
- Carter, V. and N. Rybicki, N.B. 1985. Effects of grazers and light penetration on the survival of transplants of *Vallisneria americana* Michx in the tidal Potomac River, Maryland. Aquat. Bot. 23: 197-213.
- Chamberlain, R.H., P.H. Doering, 1998a. Freshwater inflow to the Caloosahatchee Estuary and the resource-based method for evaluation. Proceedings of The Charlotte Harbor Public Conference and Technical Symposium. Charlotte Harbor National Estuary Program.
- Chamberlain, R.H., P.H. Doering, 1998b. Preliminary estimate of optimum freshwater inflow to the Caloosahatchee Estuary: A resource-based approach. Proceedings of The Charlotte Harbor Public Conference and Technical Symposium. Charlotte Harbor National Estuary Program, Tech. Report 98-02
- Chamberlain, R.H., D.E. Haunert, P.H. Doering, K.M. Haunert, J.M. Otero, and A. D. Steinman 1995. Preliminary estimate of optimum freshwater inflow to the Caloosahatchee Estuary, Florida, Department of Ecosystem Restoration, South Florida Water Management District, West Palm Beach, Fl.
- Cox, J., R. Kautz, M. MacLaughlin, and T. Gilbert. 1994. Closing the Gaps in Florida's Wildlife HabitatConservation System. Florida Game and Freshwater Fish Commission. Tallahassee. Florida
- Cummins, K.W. and Merritt, R.W. 1999. Biomonitoring and Management of North American Freshwater Wetlands, New York: Wiley & Sons, Inc.

- Day, J.W., C.A. S. Hall, W.M. Kemp, and A. Yanez-Arancibia, 1989. Estuary Ecology. John Wiley and Sons, Inc. 558 pp.
- DeGrove, B.D 1981. Caloosahatchee River watershed allocation documentation. Florida Department of Environmental Regulation, Water Quality Tech. Ser. 2(52).
- Degrove, B.D and F. Nearhoff. 1987. Water quality assessment for the Caloosahatchee River, Florida, Dept. of Environmental Regulation, Water Quality Tech. Ser. 3(19).
- Doering, P.H. and R.H. Chamberlain 1999. Water quality and the source of freshwater discharge to the Caloosahatchee Estuary, Fl. Water Resources Bulletin 35: 793-806.
- Doering, P.H., R.H. Chamberlain, K.M. Donohue, and A.D. Steinman 1999. Effect of salinity on the growth of *Vallisneria americana* Michx. from the Caloosahatchee Estuary, Florida. Florida Scientist 62(2): 89-105.
- Doering, P.H., R.H. Chamberlain, and J.M. McMunigal, 2000. Effects of simulated saltwater intrusions on the growth and survival of wild celery, *Vallisneria americana* from the Caloosahatchee Estuary (South Florida. Estuaries (submitted).
- Doering, P.H., memo to Dan Haunert dated March 22, 2000, *Vallisneria* growth algorithm, Water Research and Planning Department, South Florida Water Management District, West Palm Beach, Fl.
- Drew, D.R and Schomer, N.S. 1984. An Ecological Characterization of the Caloosahatchee River/Big Cypress Watershed. U.S. Fish Wild. Serv. FWS/OBS-82/58.2. 225 pp.
- Duever, M.J. 1984. Environmental factors controlling plant communities of the Big Cypress Swamp. In: Gleason, P.J. (Ed.), Environments of South Florida: Present and Past. II. Miami Geological Society, Miami, FL. Pp. 127-37.
- Duever, M.J., J.E. Carlson, J.F. Meeder, L.C. Duever, L.H. Gunderson, L.A. Riopelle, T.R. Alexander, R.L. Meyers and D. Spangler. 1986. The Big Cypress National Preserve. New York: National Audubon Society. 444 pp.
- Duever, M.J. 1988. Hydrologic processes for models of freshwater wetlands. In: Mitsch, William, J.M. Jorgensen and S.E. Jorgensen (Eds.), Wetlands Modeling. Amsterdam: Elsevier. pp. 9-39.
- Erwin, K. 1991. South Florida Water Management District wetland mitigation study, Volume 1. Report to the South Florida Water Management District, West Palm Beach, FL. 124 pp
- Estevez, E.D., 2000. A review and application of literature concerning freshwater flow management in riverine estuaries. Mote Marine Laboratory technical Report Number 680.
- Fan. A. and R.Burgess, 1983. Surface water availability of the Caloosahatchee basin. Tech. Memorandum. DRE-163. Department of Research and Evaluation, South Florida Water Management District, West Palm Beach, Fl. 79 pp.
- Fernald, E.A. and Purdum, E.D. 1998. Water Resources Atlas of Florida, Tallahassee: Florida State University, Institute of Science and Public Affairs.
- Fonseca, M.S. and J.S. Fisher 1986. A comparison of canopy friction and sediment movement between four species of seagrass and with reference to their ecology and restoration. Marine Ecology Progress Series 29:15-22.

- Gunter, G. 1967. Some relationships of estuaries to the fisheries of the Gulf of Mexico. In: Estuaries (Publication No. 83) (Ed: Lauff, G.H., American Association for the Advancement of Science, Washingoton, D.C. pp. 621-638.
- Gunter, G. and G.E. Hall 1962. Biological investigation of the Caloosahatchee estuary in connection with Lake Okeechobee discharges through the Caloosahatchee River. Consultant report to U.S. Corps of Engineers, Jacksonville District, Serial No. 25, 59 pp.
- Hall, C.A. 1991. Lake Okeechobee Supply-Side Management Plan. Operations and Maintenance Dept. South Florida Water Management District, West Palm Beach, Fl.
- Haller, W.T., D. L. Sutton, and W. C. Barlowe. 1974. Effects of salinity on growth of several aquatic macrophytes. Ecology 55: 891-894
- Harris, B.A., K.D. Haddad, K.A. Steidinger, and J.A. Huff 1983. Assessment of fisheries habitat: Charlotte Harbor and Lake Worth, Florida, Final Report. Florida Department of Natural Resources, Bur. of Marine Research, St. Petersburg, Fl.
- Herr, J.W. and J.E. Shaw, 1989. South Florida Water Management District ambient ground water quality. Tec. Pub. 89-1. Water quality Division, South Florida Water Management District, West Palm Beach, Fl.
- Heck, K.L., K. Able, C.Roman, and M. Fahay. 1995. Composition, abundance, biomass and production of macrofauna in a New England estuary: comparison among eelgrass meadows and other nursery habitats. Estuaries 18: 379-389.
- Haunert, D. K. and R.H. Chamberlain 1994. St. Lucie and Caloosahatchee estuary performance measures for alternative Lake Okeechobee schedules. Memo In: Special Report-Lake Okeechobee regulation schedule simulation, South Florida regional routing model. J. M. Otero and V. Floris (eds.) SFWMD, prep. For U.S. Army Corps of Engineers, Jacksonville District.
- Haunert, D.K., K.D. Konyha, P.H. Doering, R.H. Chamberlain (in review). Technical criteria to guide an adaptive management strategy for minimum flows to the Caloosahatchee Estuary, Florida. Water Research and Planning Department, South Florida Water Management District, West Palm Beach, Fl.
- Hoffacker, V.A. 1994. Caloosahatchee River submerged grass observations. W. Dexter Bender and Ass. Inc., Letter-report and Map to Chip Merriam, South Florida Water Management District, Ft. Myers, Fl.
- Irlandi, E.A., W.G. Ambrose, Jr. and B.A. Orlando 1995. Landscape ecology and the marine environment: how spatial configuration of seagrass habitat influences growth and survival of the bay scallop. Oikos 72: 307-313.
- Jassby, A.D. and Others. 1995. Isohaline position as a habitat indicator for estuarine populations. Ecol. Applications 5: 272-289.

- Kemp, W.M., W. R. Boynton, R. R. Twilley, J. C. Stevenson, and L. G. Ward. 1984. Influences of submersed vascular plants on ecological processes in upper Chesapeake Bay. pp. 367-394 in: Kennedy, V.S. (ed), The estuary as a filter. Academic Press, New York, New York.
- Killgore, K. J., R. P. Morgan II, and N. B. Rybicki. 1989. Distribution and abundance of fishes associated with submersed aquatic plants in the Potomac River. North American Journal of Fisheries Management 9:101-111.
- Konyha, K., memo dated June 26 1999. Establishing interim MFL release rules to avoid significant harm to *Vallisneria* in the Caloosahatchee Estuary, Watershed Research and Planning Department, South Florida Water Management District, West Palm Beach, Fl.
- Konyha, K,. memo dated June 29,2000.Modeling of the environmental response of *Vallisneria* (tape grass) in the Caloosahatchee Estuary to freshwater flows at the Franklin Lock & Dam (S-79, Watershed Research and Planning Department, South Florida Water Management District, West Palm Beach, Fl.
- Lubbers, L., W. R. Boynton, and W. M. Kemp. 1990. Variations in structure of estuarine fish communities in relation to abundance of submersed vascular plants. Marine Ecology Progress Series 65: 1-14.
- Mazzotti, F.E., L.A. Brandt, L.G. Pearlstrine, W.M. Kitchens, T.A. Obreza, F.C. Depkin, N.E. Morris, and C.E. Arnold. 1992. An evaluation of the regional effects of new citrus development on the ecological integrity of wildlife resources in southwest Florida. Final Report. South Florida Water Management District, West Palm Beach, FL 188 pp.
- Meyers, R.L. and J.J. Ewel, 1990. Ecosystems of Florida. Gainesville: University presses of Florida.
- Miller, T.H. 1980. Water quality aspects of the Caloosahatchee River system. Phase II, 1978-1979. Tech. Memo., South Florida Water Management District, West Palm Beach, Fl.
- Mitch, W.J. and J.G. Gosselink. 1986. Wetlands. New York: Van Nostrand Reinhold Company.
- Myers, R. and Ewel, J. 1990. Ecosystems of Florida. Orlando: University of Florida Press.
- Orth, R.J. K.L. Heck, Jr., and J. van Montfrans 1984. Faunal communities in seagrass beds: a review of the influence of plant structure and prey characteristics on predator-prey relationships. Estuaries 7: 339-350.
- Otero, J.M., J.W. Labadie, D.E. Haunert, and M.S. Daron 1995. Optimization of managed runoff to the St. Lucie Estuary. In: Water Resources Engineering, Vol. 2. W.H. Esprey, Jr. and P.G. Cobbs (eds). Proceedings First Intern. Conf., Water Resources Engineering, Amer. Soc. Civil Engineers (ASCE). 1506-1510.
- Peterson, C.H. 1982. Clam predation by whelks (*Busycon* spp.): experimental tests of the importance of prey size, prey density, and seagrass cover. Mar. Biol. 66: 159-170.
- Thayer, G.W., K.A. Bjorndal, J.C. Ogden, S.L. Williams, and J.C. Zieman 1984. Role of larger herbivores in seagrass communities. Estuaries 7: 351-376.

- Phillips, R.C and V.G. Springer 1960. A report on the hydrography, marine plants, and fishes of the Caloosahatchee River ara, Lee County, Florida. Fla. State Bd. Conserv. Marine Lab. Special Science report, No. 5, 34 pp.
- Scarlatos, P.D. 1988. Caloosahatchee Estuary Hydrodynamics. SFWMD Technical Publication No. 88-7. 39 pp.
- SFWMD, 1980. District water Use and Supply Development Plan, Volume III. Lower West Coast Planning Area. . Resource Planning Department, South Florida Water Management District, West Palm Beach, FL. April 1980
- SFWMD, 1989. Lake Okeechobee Surface Water Management and Improvement (SWIM) Plan. Part I: Water Quality and Part VIII: Public Information, Planning Department, South Florida Water Management District, West Palm Beach, FL.
- SFWMD 1992. Water supply needs and sources. Planning Department, South Florida Water Management District, West Palm Beach, FL.
- SFWMD, 1997a. Surface Water Management and Improvement (SWIM) –Update for Lake Okeechobee, Vol 1, Planning Department, South Florida Water Management District, West Palm Beach, FL.
- SFWMD,1997b. Basis of Review for Water Use. Management of Water Use Permitting Information Manual Volume III for the South Florida Water Management District, South Florida Water Management District, West Palm Beach, FL.
- SFWMD, 1999. A primer to the South Florida Water Management Model, version 3.5, Hydrologic Systems Modeling Division, South Florida Water Management District, West Palm Beach, FL. 223 pp.
- SFWMD, 2000a. District Water Management Plan, Office of Finance, South Florida Water Management District, West Palm Beach, FL. 223 pp.
- SFWMD, 2000b. Lower West Coast Water Supply Plan, Water Supply Department, Water Resources Management, South Florida Water Management District, West Palm Beach, FL.
- SFWMD, 2000c. Lower East Coast Regional Water Supply Plan, Water Supply Planning and Development Department, South Florida Water Management District, West Palm Beach, FL.
- SFWMD, 2000d. Caloosahatchee Water Management Plan, Water Supply Department, Water Resources Management, South Florida Water Management District, West Palm Beach, FL.
- SFWMD, 2000e. Minimum Flows and levels for Lake Okeechobee, the Everglades, and the Biscayne Aquifer, Water Supply Department, Water Resources Management, South Florida Water Management District, West Palm Beach, FL.
- SFWMD, 2000f. Upper East Coast Regional Water Supply Plan, Water Supply Department, Water Resources Management, South Florida Water Management District, West Palm Beach, FL.

- Twilley, R., W. Kemp, K. Staver, J. Court Stevenson, and W. Boynton. 1985. Nutrient enrichment of estuarine submersed vascular plant communities. 1. Algal growth and effects on production of plants and associated communities. Mar. Ecol. Prog. Ser. 23: 179-191.
- Twilley, R.R. and J. W. Barko. 1990. The growth of submerged macrophytes under experimental salinity and light conditions. Estuaries 13: 311 321.
- USACE 1957. General Design Memorandum, Caloosahatchee River and Control structures (Canal C-43 and lock spillway structures 77, 78, and 79) Part IV, Central Southern Florida project, USACE, Serial No. 36.
- USFWS, 1957. A preliminary survey of the fish and wildlife resources in relation to the Corps of Engineers plan of development, Caloosahatchee River-Canal C-43, Fl. U.S.G.A.S. Region 4, Atalanta, Ga.
- USACE and SFWMD, 1999. The Central and Southern Florida Flood Control Project Comprehensive Review Study. U.S Army Corps of Engineers, Jacksonville District, Jacksonville, Fl. and South Florida Water Management District, West Palm Beach, Fl.
- USACE, 2000a. Lake Okeechobee Master Water Control Manual
- USACE, 2000b. Lake Okeechobee Regulation Schedule Study: Final Environmental Impact Statement, U.S. Army Corps of Engineers, Jacksonville, District.
- USEPA, 1987. Estuary Program Primer, Environmental Protection Agency, Office of Marine and Estuarine Protection, Washington, D.C.
- Wagner, C.M. and H.M. Austin 1999. Correspondence between environmental gradients and summer littoral fish assemblages in low salinity reaches of the Chesapeake Bay, USA. Mar. Ecol. Prog Ser. 177: 197-212.
- Zieman, J.C., and R.T Zieman. 1989. The ecology of the seagrass meadows of the west coast of Florida: a community profile. U. S. Fish and Wildlife Service Biological Report 85(7.25). 155 pp.